

A Note on Partial List Coloring

Moharram N. Iradmusa¹

Department of Mathematical Sciences

Shahid Beheshti University

P.O. Box 19834, Tehran, Iran

iradmusa@yahoo.com

Abstract

Let G be a simple graph with n vertices and list chromatic number $\chi_\ell(G) = \chi_\ell$. Suppose that $0 \leq t \leq \chi_\ell$ and each vertex of G is assigned a list of t colors. Albertson, Grossman and Haas [1] conjectured that at least $\frac{tn}{\chi_\ell}$ vertices of G can be colored from these lists.

In this paper we find some new results in partial list coloring which help us to show that the conjecture is true for at least half of the numbers of the set $\{1, 2, \dots, \chi_\ell(G) - 1\}$. In addition we introduce a new related conjecture and finally we present some results about this conjecture.

Key words: List chromatic number, List coloring, List assignment.

Subject classification: 05C15 .

1 Introduction and Preliminaries

Throughout this paper we only consider finite and simple graphs. Let G be a simple graph with vertex set $V(G)$, $|V(G)| = n$ and edge set $E(G)$. For each vertex $v \in V(G)$ let $\mathcal{L}(v)$ be a list of allowed colors assigned to v . The collection of all lists is called a *list assignment* and denoted by $\mathcal{L}$. In special case we have *t-list assignment* if $|\mathcal{L}(v)| = t$ for all $v \in V(G)$. Also we call $R(\mathcal{L}) = \bigcup_{v \in V(G)} \mathcal{L}(v)$ the *color list* of $\mathcal{L}$.

The graph G is called $\mathcal{L}$ -*list colorable* if there is a coloring $c : V(G) \rightarrow R(\mathcal{L})$ such that $c(v) \neq c(u)$ for all $uv \in E(G)$ and $c(v) \in \mathcal{L}(v)$ for all $v \in V(G)$. Furthermore G is k -*choosable* if it is $\mathcal{L}$ -list colorable for every k -list assignment $\mathcal{L}$. The *list chromatic number* or *choice number*, $\chi_\ell(G) = \chi_\ell$ is the smallest number k such that G is k -choosable. List coloring was introduced independently by Vising [6] and by Erdos, Rubin and Taylor [3].

In addition, let $\lambda_{\mathcal{L}}(G)$ be the maximum number of vertices of G which are colorable with respect to the list assignment $\mathcal{L}$. Define $\lambda_t(G) = \min \lambda_{\mathcal{L}}(G)$, where the minimum is taken over all t -list assignments $\mathcal{L}$. Clearly if $t \geq \chi_\ell(G)$ then $\lambda_t = n$.

¹This paper is partially supported by Shahid Beheshti University.

Thus it is interesting to know about λ_t when $t < \chi_\ell$.

Conjecture 1. (Albertson, Grossman and Haas [1]) Let G be a graph with n vertices and list chromatic number $\chi_\ell(G) = \chi_\ell$. Then, for any $0 \leq t \leq \chi_\ell$ we have $\lambda_t \geq \frac{tn}{\chi_\ell}$.

We call this conjecture *AGH conjecture*. The conjecture is clearly correct for $t = 0$, $t = 1$ and $t = \chi_\ell$. Albertson et. al. [1] proved the following theorem:

Theorem 1. Let $0 \leq t \leq \chi_\ell$. Then $\lambda_t \geq (1 - (1 - \frac{1}{\chi})^t)n$ and this number is asymptotically best possible.

Furthermore, Chappell [2] found the lower bound $\frac{6}{7} \frac{tn}{\chi_\ell}$ for λ_t for all t between 0 and χ_ℓ . In addition Haas et. al. [4] showed that the conjecture is true when $t|\chi_\ell$. For more information see also [5] and [7].

In this paper we find some new results which help us to show that *AGH conjecture* is true for at least half of the numbers of the set $\{1, 2, \dots, \chi_\ell(G) - 1\}$. Finally we introduce a new conjecture and present some results about this conjecture.

2 Main Results

In the first theorem of this section we prove a fundamental inequality about the numbers λ_t where $0 \leq t \leq \chi_\ell$.

Theorem 2.(Triangle Inequality) Let $1 \leq r, s \leq \chi_\ell$. Then,

$$\lambda_r + \lambda_s \geq \lambda_{r+s}.$$

Proof. Let $\mathcal{L}_r$ and $\mathcal{L}_s$ be r -list and s -list assignments of G respectively, such that $\lambda_{\mathcal{L}_r} = \lambda_r$ and $\lambda_{\mathcal{L}_s} = \lambda_s$. Also suppose that $R(\mathcal{L}_r) \cap R(\mathcal{L}_s) = \emptyset$. We define an $(r + s)$ -list assignment $\mathcal{L}_{r+s}$ on $V(G)$ with

$$\mathcal{L}_{r+s}(v) = \mathcal{L}_r(v) \cup \mathcal{L}_s(v).$$

Then we have $\lambda_{r+s} \leq \lambda_{\mathcal{L}_{r+s}}$. Now suppose that $c : V(G) \rightarrow R(\mathcal{L}_{r+s})$ is a partial list coloring of G such that $\lambda_{\mathcal{L}_{r+s}}$ vertices of G are colored. Then we can divide colored vertices of G to the sets $R = \{v \in V(G) | c(v) \in \mathcal{L}_r(v)\}$ and $S = \{v \in V(G) | c(v) \in \mathcal{L}_s(v)\}$. This partition results $|R| \leq \lambda_{\mathcal{L}_r}$ and $|S| \leq \lambda_{\mathcal{L}_s}$. Finally,

$$\lambda_{r+s} \leq \lambda_{\mathcal{L}_{r+s}} = |R| + |S| \leq \lambda_{\mathcal{L}_r} + \lambda_{\mathcal{L}_s} = \lambda_r + \lambda_s.$$

■

Corollary 3. Let G be a graph with list chromatic number $\chi_\ell(G) = \chi_\ell$.

1. If $1 \leq r_i \leq \chi_\ell (1 \leq i \leq k)$, then $\sum_{i=1}^k \lambda_{r_i} \geq \lambda_{\sum_{i=1}^k r_i}$.

In particular case, if $r_1 = r_2 = \dots = r_k = r$, then we have $k\lambda_r \geq \lambda_{kr}$.

2. If $1 \leq r, s \leq \chi_\ell$ and $r|s$, then $\frac{s}{r}\lambda_r \geq \lambda_s$ or $\frac{\lambda_r}{r} \geq \frac{\lambda_s}{s}$.

The next Corollary was proved in [4]. Here we give another proof.

Corollary 4. If $t|\chi_\ell(G)$, then $\lambda_t \geq \frac{tn}{\chi_\ell}$.

Proof. In part two of Corollary 3 set $s = \chi_\ell$ and use $\lambda_{\chi_\ell} = |V(G)|$. ■

As a result of Theorem 2 we can prove that *AGH conjecture* is true for at least half of the numbers of the set $\{1, 2, \dots, \chi_\ell(G) - 1\}$.

Corollary 5. Let $1 \leq r \leq \chi_\ell = s$. Then *AGH conjecture* is true for r or $s - r$. In other words, at least one of the following inequalities is true:

$$\lambda_r \geq \frac{rn}{\chi_\ell} \quad , \quad \lambda_{s-r} \geq \frac{(s-r)n}{\chi_\ell}.$$

Proof. Suppose that $\lambda_r < \frac{rn}{\chi_\ell}$ and $\lambda_{s-r} < \frac{(s-r)n}{\chi_\ell}$. Then

$$\lambda_r + \lambda_{s-r} < \frac{rn}{\chi_\ell} + \frac{(s-r)n}{\chi_\ell} = n.$$

Furthermore from Theorem 2 we have $\lambda_r + \lambda_{s-r} \geq \lambda_{r+s-r} = \lambda_s$.

So we conclude $\lambda_{\chi_\ell} < n$ that is a contradiction. ■

Similarly we can prove the following result:

Corollary 6. Let $1 \leq r \leq \chi_\ell$ and $\chi_\ell = kr + r_0$ where $0 \leq r_0 < r$. Then *AGH conjecture* is true for r or r_0 .

The next Corollary was proved in [4]. Here we prove it as a result of Corollary 3.

Corollary 7. If $1 \leq r \leq \chi_\ell = s$ then

$$\lambda_r \geq \frac{n}{\lceil \frac{s}{r} \rceil}.$$

Proof. From Corollary 3 we have

$$\lceil \frac{s}{r} \rceil \lambda_r \geq \lambda_{r \lceil \frac{s}{r} \rceil}.$$

Also from $r \lceil \frac{s}{r} \rceil \geq s$ we have $\lambda_{r \lceil \frac{s}{r} \rceil} = n$ which completes the proof. ■

In the next corollary we generalize the Corollaries 4 and 6.

Corollary 8. Let $1 \leq r \leq s \leq \chi_\ell$.

1. If $r|s$ and *AGH conjecture* is true for s , then it is true for r .
2. Suppose that $s = kr + r_0$ and $0 \leq r_0 < r$. If *AGH conjecture* is true for s , then it is true for r or r_0 .

Here we give a new conjecture that is a generalization of *AGH conjecture*:

Conjecture 2. Let G be a graph of order n with list chromatic number χ_ℓ and $1 \leq r \leq s \leq \chi_\ell$. Then

$$\frac{\lambda_r}{r} \geq \frac{\lambda_s}{s}.$$

Remark.

1. Part two of the Corollary 3 shows that Conjecture 2 is true when $r|s$.
2. If $s = \chi_\ell$, then $\lambda_s = n$. So *AGH conjecture* is the special case of Conjecture 2.
3. Suppose that $r = 1$. So $r|s$ and we have $\lambda_s \leq s\lambda_1 = s\alpha(G)$ where $\alpha(G)$ is the stability number of G .

In the next theorems we show some results about the new conjecture. We need some definitions which are given before the theorems.

Definition 1. Let G be a graph of order n with list chromatic number χ_ℓ . Then $Td(G)$ is the set of all ordered pairs (r, s) , such that $r \leq s$ and $\frac{\lambda_r}{r} \geq \frac{\lambda_s}{s}$.

Theorem 9. Let G be a graph of order n with list chromatic number χ_ℓ and $1 \leq r \leq s \leq \chi_\ell$.

1. We have $(r, s) \in Td(G)$ or $(s - r, s) \in Td(G)$. So Conjecture 2 is true for at least half of the elements of the set $Td(G)$.
2. If $s = kr + r_0$ and $0 \leq r_0 < r$, then $(r, s) \in Td(G)$ or $(r_0, s) \in Td(G)$.
3. We have

$$\lceil \frac{s}{r} \rceil \lambda_r \geq \lambda_s.$$

In other word $\alpha \frac{\lambda_r}{r} \geq \frac{\lambda_s}{s}$, where $\alpha = \frac{r}{s} \lceil \frac{s}{r} \rceil$.

4. The set $Td(G)$ is an order relation that is defined on the set $\{1, 2, \dots, \chi_\ell(G)\}$.

Proof. We can prove this theorem similar to the Corollaries 5,6 and 7. ■

Definition 2. Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be two list assignments of G . We say $\mathcal{L}_1 \subseteq \mathcal{L}_2$ if $\mathcal{L}_1(v) \subseteq \mathcal{L}_2(v)$ for all $v \in V(G)$.

Definition 3. Let H be a subgraph of the graph G , $\mathcal{L}'$ be a list assignment of H and $\mathcal{L}$ be a list assignment of G . We say $\mathcal{L}'$ is a *restriction of $\mathcal{L}$ to H* if $\mathcal{L}'(v) = \mathcal{L}(v)$ for all $v \in V(H)$ and we show that by $\mathcal{L}' = \mathcal{L} \upharpoonright_H$.

The next lemma is an straightforward result of the definition 3.

Lemma 10. Let H be an induced subgraph of the graph G , $\mathcal{L}$ be a list assignment of G and $\mathcal{L}' = \mathcal{L} \upharpoonright_H$. Then $\lambda_{\mathcal{L}}(G) \geq \lambda_{\mathcal{L}'}(H)$.

Theorem 11. Let H be an induced subgraph of the graph G and $1 \leq t \leq \chi_{\ell}(G)$. Then $\lambda_t(G) \geq \lambda_t(H)$.

Proof. Suppose that $\mathcal{L}$ is a t -list assignment of G such that $\lambda_{\mathcal{L}} = \lambda_t(G)$ and $\mathcal{L}' = \mathcal{L} \upharpoonright_H$. So $\lambda_t(G) = \lambda_{\mathcal{L}}(G) \geq \lambda_{\mathcal{L}'}(H) \geq \lambda_t(H)$. ■

Now we prove the following theorem for the conjecture 2, that is similar to Theorem 1.

Theorem 12. Let G be a graph of order n with list chromatic number χ_{ℓ} and $1 \leq r \leq s \leq \chi_{\ell}$. Also suppose that $\mathcal{L}_s$ is an s -list assignment of G with $\mathcal{L}_s(v) = \{1, 2, \dots, s\}$ for all $v \in V(G)$. Then

$$\lambda_r \geq (1 - (1 - \frac{1}{s})^r) \lambda_{\mathcal{L}_s}.$$

Proof. We can color at most $\lambda_{\mathcal{L}_s}$ vertices of G from the list color $\{1, 2, \dots, s\}$. Let H be a subgraph of G induced by $\lambda_{\mathcal{L}_s}$ colored vertices. At first we show that $\chi(H) = s$. Suppose that $\chi(H) \leq s - 1$. So there is a coloring of G by at most $s - 1$ colors of $\{1, 2, \dots, s\}$. Thus we can color at least one vertex of $V(G) - V(H)$ by the remaining colors such that the coloring remain proper which contradict by the definition of H . So $\chi(H) = s$. Now from Theorem 1 we have

$$\lambda_r(H) \geq (1 - (1 - \frac{1}{\chi(H)})^r) |V(H)| = (1 - (1 - \frac{1}{s})^r) \lambda_{\mathcal{L}_s}.$$

Also from Theorem 11 we have $\lambda_r(G) \geq \lambda_r(H)$. So $\lambda_r(G) \geq (1 - (1 - \frac{1}{s})^r) \lambda_{\mathcal{L}_s}$. ■

Corollary 13. By assumptions of Theorem 12 and by use of $\lambda_{\mathcal{L}_s} \geq \lambda_s$, we have

$$\lambda_r \geq (1 - (1 - \frac{1}{s})^r) \lambda_s,$$

that is similar to the result of Theorem 1.

The last theorem shows the relation between partial list coloring of a graph and its subgraphs.

Theorem 14. Let G be a graph of order n with list chromatic number χ_{ℓ} and $1 \leq t \leq \chi_{\ell}$.

(1) Suppose that H is a subgraph of G of maximum order such that $\chi_{\ell}(H) = t$. Then, $\lambda_t(G) \geq |H|$.

(2) Suppose that $\lambda_t = \lambda_{\mathcal{L}_t}$ where $\mathcal{L}_t$ is a t -list assignment of G . If H is a subgraph of G of order λ_t , induced by colored vertices in the partial coloring of G by $\mathcal{L}_t$, then $\chi_{\ell}(H) \geq t$.

Proof. (1) Suppose that $\lambda_t(G) \leq |H| - 1$ and $\lambda_t(G) = \lambda_{\mathcal{L}_t}(G)$ where $\mathcal{L}_t$ is a t -list assignment of G . Thus in the coloring of G from the lists of $\mathcal{L}_t$, we can color less than $|H|$ vertices. But $\chi_\ell(H) = t$. So by restriction of $\mathcal{L}_t$ to H , we can color all vertices of H that is a partial list coloring of G and this contradicts with $\lambda_{\mathcal{L}_t}(G) < |H|$. So $\lambda_t(G) \geq |H|$.

(2) Suppose that $\chi_\ell(H) = s \leq t - 1$. So for any s -list assignment $\mathcal{L}_s$ of H , there is a proper coloring of H by the lists of $\mathcal{L}_s$. Also consider the t -list assignment $\mathcal{L}_t$ and select arbitrary vertex x of $G - H$. Because x is uncolored, so each color of $\mathcal{L}_t(x)$ must be appeared on at least one vertex of $N(x)$. Now define an s -list assignment $\mathcal{L}'$ such that $\mathcal{L}' \subseteq \mathcal{L}$ and $c \notin \mathcal{L}'(v)$ for all $v \in V(G)$. So we can color the vertices of H by lists of $\mathcal{L}'$. Thus by adding $t - s$ removed colors to each list we can color the vertex x by the color c . So we can color $|H| + 1$ vertices of G by lists of $\mathcal{L}_t$ that is a contradiction. So $\chi_\ell(H) \geq t$. ■

Acknowledgment

We would like to thank Professor Hossein Hajiabolhassan for his useful comments.

References

- [1] M. Albertson, S. Grossman and R. Haas, *Partial list colouring*, Discrete Math., **214**(2000), pp. 235-240.
- [2] G. Chappell, *A lower bound for partial list colouring*, J. Graph Theory, **32**(1999), pp. 390-393.
- [3] P. Erdos, A. Rubin and H. Taylor, *Choosability in graphs*, Proc. West Coast Conf. On Combinatorics, Graph Theory and Computing, Congressus Numerantium XXVI, (1979), pp. 125-127.
- [4] R. Haas, D. Hanson and G. MacGillivray, *Bounds for partial list colourings*, Ars Combin., **67**(2003), pp. 27-31.
- [5] Jeannette C.M Janssen, *A partial solution of a partial list colouring problem*,
- [6] V. Vizing, *Colouring the vertices of a graph in prescribed colors*(russian), Diskret Analiz., **29**(1976), pp. 3-10.
- [7] M. Voigt, *Algorithmic aspects of partial list colourings*, Combin. Probab. and Comput., **9**(2000), pp. 375-380.